

PREDICTING PNEUMONIA IN PAEDIATRIC ACUTE FEBRILE RESPIRATORY ILLNESS

DR. FIONA Y. Y. CHAN

ACCIDENT AND EMERGENCY DEPARTMENT, TUEN MUN HOSPITAL

COAUTHORS: DR. CF TSE¹, DR. KM POON², DR. CT LUI³

¹DEPARTMENT OF ACCIDENT AND EMERGENCY, PRINCESS MARGARET HOSPITAL

²DEPARTMENT OF ACCIDENT AND EMERGENCY, POK OI HOSPITAL, HONG KONG

³DEPARTMENT OF ACCIDENT AND EMERGENCY, TUEN MUN HOSPITAL

INTRODUCTION

- Pneumonia
 - 932 per 100,000 population of 0-5 years old admitted to hospital in HK
 - Kills 1.6 million paediatric patients per year
 - 16% of all deaths in children <5 years old

INTRODUCTION



+



= ?

fever

cough

URTI ? Pneumonia? Bronchitis?

INTRODUCTION

- To take CXR, or not to take, that is the question.
 - Hamlet ver. 2.0, SSEM 2017



PREDICTORS IN THE PAST

- World Health Organization (WHO) 1991
- **cough, tachypnea and chest indrawing** for pneumonia diagnosis
- Validated in various studies
- poor sensitivity for diagnosing pneumonia in a US-based paediatric emergency department (2012)

PROPOSED PREDICTORS (2004-2011)

	Al-Dabbagh et al (2004)	Lynch et al (2004)	Bilkis prediction rule (2010)	Bilkis simpler rule (2010)	Neuman et al (2011)
Study Design	Case series	Prospective cohort study	Prospective cohort study	Prospective cohort study	Prospective cohort study
Clinical Predictors	nasal flaring, tachypnea, chest indrawing and crackles parallel testing for sensitivity	fever + decreased breath sounds, crackles or tachypnea	Grunting, cough, rales, decreased breath sounds, vomiting	Decreased breath sounds, rales, tachypnea or any combination of the 3	chest pain, focal rales, duration of fever, and oximetry levels at triage
	Temperature >38°C and grunting tested serially for specificity				SpO ₂ ≤ 92% strongest predictor of pneumonia in <5 yo

BACKGROUND

- Our aim:
- evaluate **independent determinant factors** in prediction of pneumonia
- establish a **clinical prediction rule** for pneumonia in paediatric patients with acute febrile respiratory illness

METHOD

STUDY DESIGN AND SETTING

- Multicentre prospective cohort study
- TMH, PMH, POH
- September 2016 to May 2017
- 1 month to 6 years old

METHOD

INCLUSION CRITERIA

- acute symptom onset within **10 days**
- fever $\geq 38^{\circ}\text{C}$ **within 24 hours** at attendance at the A&E
- body temperature in triage, ambulance and self-reported; either oral, rectal or tympanic temperature
- **respiratory symptoms** (cough, dyspnea, wheezing or added sounds during respiration)

METHOD

EXCLUSION CRITERIA


- hypoxemic (SpO₂ of \leq 94%)
- obvious respiratory distress or requiring O₂ supplement
- immunocompromised
- chronic lung disease

OUTCOME

- Pneumonia
- 1. new onset radiological lung infiltrate in chest radiograph
- 2. re-attendance to any emergency departments in Hong Kong within 7 days and diagnosed pneumonia

RESULTS

PATIENT ENROLMENT – SPLIT SAMPLE METHOD



RESULTS

DERIVATION GROUP

- **Male** predominance
- Median age **26 months**

RESULTS


	All (n=484)	Positive outcome (n=48)	Negative outcome (n=436)	p value
Symptoms				
Peak temperature within 24 hours, °C [median ± IQR]	39 (38.5–39.8)	39 (38.7–39.9)	39 (38.5–39.8)	0.923
Peak temperature within 24 hours ≥ 39°C	295 (61.1%)	33 (68.8%)	262 (60.1%)	0.243
Fever duration, days [mean ± SD]	2.3 ± 1.8	3.7 ± 2.7	2.2 ± 1.6	<0.001
Fever duration ≥ 3 days	155 (32.1%)	31 (64.6%)	124 (28.4%)	<0.001
Fever duration ≥ 5 days	53 (11%)	16 (33.3%)	37 (8.5%)	<0.001
Fever duration ≥ 7 days	13 (2.7%)	5 (10.4%)	8 (1.8%)	0.005
Chills	85 (17.6%)	14 (29.2%)	71 (16.3%)	0.026
Cough	477 (98.8%)	46 (95.8%)	431 (98.9%)	0.147
Purulent sputum	23 (4.8%)	0 (0%)	23 (5.3%)	0.152
Noted dyspneic by parents	41 (8.5%)	10 (20.8%)	31 (7.1%)	0.001
Noisy breathing	40 (8.3%)	3 (6.3%)	37 (8.5%)	0.785
Sore throat	132 (27.3%)	13 (27.1%)	119 (27.3%)	0.975
Nasal blockage / discharge	370 (76.6%)	32 (66.7%)	338 (77.5%)	0.093
Poor feeding	134 (27.7%)	13 (27.1%)	121 (27.8%)	0.922
Travel history within 2 weeks	59 (12.2%)	8 (16.7%)	51 (11.7%)	0.318
Hospitalized within 7 days	7 (1.4%)	0 (0%)	7 (1.6%)	1.000

RESULTS

	All (n=484)	Positive outcome (n=48)	Negative outcome (n=436)	p value
Physical examination				
Toxic appearance	10 (2.1%)	3 (6.3%)	7 (1.6%)	0.067
Reduced breath sounds	1 (0.2%)	0 (0%)	1 (0.2%)	1.000
Crackles	16 (3.3%)	5 (10.4%)	11 (2.5%)	0.015
Wheeze on examination	14 (2.9%)	1 (2.1%)	13 (3%)	1.000
Past history				
Past history of pneumonia	49 (10.1%)	4 (8.3%)	45 (10.3%)	0.805
Asthma	18 (3.7%)	4 (8.3%)	14 (3.2%)	0.757
Presenting vital signs				
Pulse rate, beats per min [mean ± SD]	138 ± 26	136 ± 23	139 ± 26	0.186
Tachycardia (99 th percentile)	254 (52.6%)	27 (56.3%)	227 (52.1%)	0.582
Tachycardia (99 th percentile, adjusted for fever)	139 (28.8%)	15 (31.3%)	124 (28.4%)	0.683
Respiratory rate, breaths per min [median ± IQR]	20 (18–25)	20 (18–24.5)	20 (18–25)	0.967
Tachypnea (99th percentile)	11 (2.3%)	4 (8.3%)	7 (1.6%)	0.017
SpO ₂ , % [median ± IQR]	98 (97–99)	97.5 (96–98)	98 (97–99)	0.007
SpO₂ ≤ 96%	92 (19%)	19 (39.6%)	73 (16.7%)	<0.001
Temperature, °C [median ± IQR]	38.1 (37.2–38.8)	38.1 (37.4–38.9)	38.1 (37.2–38.8)	0.489
Temperature ≥ 39°C	82 (17%)	6 (12.5%)	76 (17.4%)	0.387

RESULTS

Association between SpO₂ and probability of pneumonia in pediatric patients



PAEDIATRIC AFRI RULE (WITH RESPIRATORY RATE)

	Log OR	Wald	Odds ratio (95% CI)	p value	Weight
Duration of fever		47.37		<0.001	
< 3 days			1		
3-4 days	0.811	7.74	2.25 (1.27 - 3.98)	0.005	2
5-6 days	1.862	32.60	6.44 (3.4 - 12.2)	<0.001	4
≥ 7 days	2.18	25.26	8.84 (3.78 - 20.69)	<0.001	5
Chills	0.696	6.93	2.01 (1.2 - 3.37)	0.008	2
Nasal symptoms	-0.767	9.17	0.46 (0.28 - 0.76)	0.002	-2
Abnormal chest examination	1.183	10.66	3.27 (1.61 - 6.65)	0.001	3
Tachypnea > 99th percentile	1.384	5.22	3.99 (1.22 - 13.1)	0.022	3

Backward stepwise method by likelihood ratio

Hosmer-and-lemeshow goodness-of-fit test p=0.405

AUROC of predicted probabilities 0.762 (95% CI 0.688-0.830)

PAEDIATRIC AFRI RULE (WITH SPO₂)

	Log OR	Wald	Odds ratio (95% CI)	p value	Weight
Duration of fever		45.467		<0.001	
< 3 days			1		
3-4 days	0.799	7.562	2.22 (1.26 - 3.93)	0.006	2
5-6 days	1.85	32.416	6.36 (3.36 - 12.02)	<0.001	4
≥ 7 days	2.102	23.027	8.18 (3.47 - 19.3)	<0.001	5
Chills	0.705	7.181	2.03 (1.21 - 3.39)	0.007	2
Nasal symptoms	-0.724	8.279	0.49 (0.3 - 0.79)	0.004	-2
Abnormal chest examination	1.156	9.998	3.18 (1.55 - 6.5)	0.002	3
SpO₂ ≤ 96%	0.952	2.961	2.59 (1.94 - 3.69)	0.035	3

Backward stepwise method by likelihood ratio


Hosmer-and-lemeshow goodness-of-fit test p=0.851

AUROC of predicted probabilities 0.761 (95% CI 0.683-0.832)

PROPOSED PAEDIATRIC AFRI (PAFRI) RULE

Predictors	Weight
Duration of fever	
< 3 days	0
3-4 days	2
5-6 days	4
≥ 7 days	5
Chills (observed chills by parents)	2
Nasal symptoms (nasal discharge, nasal blockage)	-2
Abnormal chest examination (any of reduced breath sound, rales, and wheeze)	3
SpO ₂ ≤ 96% or tachypnea	3

COMPARISON BETWEEN SCORES



AUROC:

PAFRI 0.733 (95% CI 0.653-0.813; $p < 0.001$),
Bilkis decision rule 0.600 (95% CI 0.510-0.691;
 $p = 0.03$),

Bilkis simpler rule 0.579 (95% CI 0.481-0.678;
 $p = 0.086$)

Non-parametric Comparison of AUROC:

PAFRI vs Bilkis decision rule $p < 0.001$;

PAFRI vs Bilkis simpler rule $p < 0.001$.

DIAGNOSTIC CHARACTERISTICS OF PAFRI RULE AT VARIOUS CUT-OFFS (IN DERIVATION DATA SET)

	Outcome	PAFRI ≥ 0	PAFRI ≥ 1	PAFRI ≥ 2	PAFRI ≥ 3	PAFRI ≥ 4	PAFRI ≥ 5
Positive test	Positive	44	37	34	27	17	14
	Negative	269	150	113	68	33	25
Negative test	Positive	4	11	14	21	31	34
	Negative	167	286	323	238	403	411
Sensitivity		91.7 (79.1-97.3)	77.1 (62.3-87.5)	70.8 (55.7-82.6)	56.3 (41.3-70.2)	35.4 (22.6-50.6)	29.2 (17.4-44.3)
Specificity		38.3 (33.7-43.1)	65.6 (60.9-70)	74.1 (69.7-78.1)	77.8 (72.6-82.2)	92.4 (89.4-94.7)	94.3 (91.5-96.2)
PPV		14.1 (10.5-18.5)	19.8 (14.5-26.4)	23.1 (16.8-30.9)	28.4 (19.9-38.7)	34 (21.6-48.9)	35.9 (21.7-52.8)
NPV		97.7 (93.7-99.2)	96.3 (93.3-98)	95.8 (93-97.6)	91.9 (87.7-94.8)	92.9 (89.9-95)	92.4 (89.4-94.6)
PLR		1.49 (1.33-1.66)	2.24 (1.83-2.74)	2.73 (2.15-3.48)	2.53 (1.83-3.51)	4.68 (2.83-7.74)	5.09 (2.84-9.11)
NLR		0.22 (0.08-0.56)	0.35 (0.21-0.59)	0.39 (0.25-0.61)	0.56 (0.41-0.78)	0.7 (0.57-0.86)	0.75 (0.63-0.9)
Youden's J		0.30	0.43	0.45	0.34	0.28	0.24

DISCUSSION

- Tachypnea VS SpO₂? similar accuracy.
- Pulse oximetry more objective, less observer dependent, quickly obtained
- RR not accurate in crying or irritable children in emergency setting.
- Pulse oximetry may not be available in other clinical settings e.g. primary care clinics.

DISCUSSION

- Fever:
 - Duration
 - Temperature at triage 艦
 - Peak Temperature 艦
- Nasal Symptoms VS Sore throat 艦

DISCUSSION – HOW TO USE THE RULE

- Clinical uses of the PAFRI:
- PAFRI ≥ 1 \rightarrow assist emergency physicians to consider ordering an CXR/help consider referral for a CXR.
- PAFRI ≥ 4 or above has $>90\%$ specificity
- Triage for early assessment with positive prediction by PAFRI
- Consider triage-initiated chest radiograph

LIMITATIONS

- Verification bias – CXR not taken in every patient
- Not done throughout the whole year – flu surge was not taken into account of in the study

CONCLUSION

- A clinical prediction rule for CXR confirmed paediatric pneumonia has been established using clinical history, signs and symptoms.
- The PAFRI rule has achieved satisfactory accuracy in performance in predicting pneumonia compared to other previous studies.
- Different cutoff points can be employed in different settings.
- To be further validated by a primary care population.

REFERENCES

1. Black RE, Cousens S, Johnson HL, Lawn JE, Rudan I, Bassani DG, et al. Global, regional, and national causes of child mortality in 2008: a systematic analysis. *Lancet*. 2010;375:1969–87.
2. Liu L, Johnson HL, Cousens S, Perin J, Scott S, Lawn JE, et al. Child Health Epidemiology Reference Group of WHO and UNICEF. Global, regional, and national causes of child mortality: an updated systematic analysis for 2010 with time trends since 2000. *Lancet*. 2012 Jun 9;379(9832):2151-61. doi: 10.1016/S0140-6736(12)60560-1. Epub 2012 May 11.
3. Pneumonia Fact Sheet, Media Centre, Sept 2016 (<http://www.who.int/mediacentre/factsheets/fs331/en/>) accessed 20 August 2017.
4. DC Lung, DSY Lam, E Chan, KC Chan, SSS Chiu, PL Ho, et al. Practice Recommendations for Management of Community Acquired Pneumonia in Children. *HK J Paediatr (new series)* 2016;21:178-193
5. Phelan PD, Olinsky A. Clinical patterns of acute respiratory infection. In: *Respiratory illness in children, 3rd ed*: Blackwell Scientific Publications; 1990.
6. Murphy, C.G., van de Pol, A.C., Harper, M.B., Bachur, R.G. Clinical predictors of occult pneumonia in the febrile child. *Acad Emerg Med*. 2007;14:243–249.
7. Programme for the Control of Acute Respiratory Infections, WHO. Technical bases for the WHO recommendations on the management of pneumonia in children at first-level health facilities. Geneva, 1991. (http://apps.who.int/iris/bitstream/10665/61199/1/WHO_ARI_91.20.pdf) accessed 20 August, 2017.
8. Campbell H, Byass P, Lamont AC, Forgie IM, O'Neill KP, Lloyd-Evans N et al. Assessment of clinical criteria for identification of severe acute lower respiratory tract infections in children. *Lancet*. 1989 Feb 11;1(8633):297-9.
9. Redd SC, Vreuls R, Metsing M, Mohobane PH, Patrick E, Moteetee M. Clinical signs of pneumonia in children attending a hospital outpatient department in Lesotho. *Bulletin of the World Health Organization*. 1994;72(1):113-118
10. Cherian T, John TJ, Simoes E, Steinhoff MC, John M. Evaluation of simple clinical signs for the diagnosis of acute lower respiratory tract infection. *Lancet*. 1988 Jul 16;2(8603):125-8.
11. Shann F, Hart K, Thomas D. Acute lower respiratory tract infections in children: possible criteria for selection of patients for antibiotic therapy and hospital admission. 1984. *Bull World Health Organ*. 2003;81(4):301-5. Epub 2003 May 16.

REFERENCES

12. Mulholland EK, Simoes EA, Costales MO, McGrath EJ, Manalac EM, Gove S. Standardized diagnosis of pneumonia in developing countries. *Pediatr Infect Dis J*. 1992 Feb;11(2):77-81.
13. Rambaud-Althaus C, Althaus F, Genton B, D'Acremont V. Clinical features for diagnosis of pneumonia in children younger than 5 years: a systematic review and meta-analysis *The Lancet Infectious Diseases* , Volume 15 , Issue 4 , 439 – 450
14. Wingerter, Sarah L, Bachur, Richard G; Monuteaux, Michael C; Neuman, Mark I. Application of the World Health Organization Criteria to Predict Radiographic Pneumonia in a US-based Pediatric Emergency Department *Pediatric Infectious Disease Journal*. 31(6):561-564, June 2012.
15. Al-Dabbagh SA, Al-Zubaidi SN. The validity of clinical criteria in predicting pneumonia among children under five years of age. *J Fam Community Med* 2004;11:11-6
16. Lynch T, Platt R, Gouin S, Larson C, Patenaude Y. Can we predict which children with clinically suspected pneumonia will have the presence of focal infiltrates on chest radiographs? *Pediatrics*. 2004;113(3 pt 1).
17. Neuman MI, Monuteaux MC, Scully KJ, Bachur RG., Prediction of Pneumonia in a Pediatric Emergency Department. *Pediatrics*. 2011 Aug;128(2):246-53. doi: 10.1542/peds.2010-3367. Epub 2011 Jul 11.
18. Bilkis MD, Gorgal N, Carbone M, Vazquez M, Albanese P, Branda MC, et al. Validation and development of a clinical prediction rule in clinically suspected community-acquired pneumonia. *Pediatr Emerg Care*. 2010;26(6):399 – 405
19. Fleming S, Thompson M, Stevens R, Heneghan C, Plüddemann A, Maconochie I et al. Normal ranges of heart rate and respiratory rate in children from birth to 18 years of age: a systematic review of observational studies. *The Lancet*, Volume 377 , Issue 9770 , 1011 - 1018
20. Davies P, Maconochie I. The relationship between body temperature, heart rate and respiratory rate in children *Emergency Med J*. 2009 Sep;26(9):641-3. doi: 10.1136/emj.2008.061598.
21. Gentile A, Bardach A, Ciapponi A, Garcia-Marti S, Aruj P, Glujovsky D et al, Epidemiology of community-acquired pneumonia in children of Latin America and the Caribbean: a systematic review and meta-analysis, *International Journal of Infectious Diseases*, Volume 16, Issue 1, 2012, Pages e5-e15, ISSN 1201-9712
22. Wahr JA, T K, Diab M. Pulse Oximetry. *Respiratory Care Clinics of North America*. 1995;1(1):77–105
23. Verhoeven D, Teijaro J, Farber DL. Pulse-oximetry accurately predicts lung pathology and the immune response during influenza infection. *Virology*. 2009;390(2):151-156. doi:10.1016/j.virol.2009.05.004.

Thank you!